



December 2024: Southwest Climate Outlook

Stacie Reece January 6, 2025



https://climas.arizona.edu/

The Southwest Climate Outlook is published by the Climate Assessment for the Southwest (CLIMAS), with support from University of Arizona Cooperative Extension, and the New

Mexico State Climate office.

Questions/Contact: Stacie Reece, sreece@arizona.edu

Precipitation and Temperature

November precipitation was generally above normal for New Mexico and below normal for Arizona. Parts of eastern New Mexico received record precipitation while for parts of western Arizona it was the driest November on record. Southwest - Precipitation November 2024, Percentile



November temperatures were near normal or below normal for much of Arizona and New Mexico. Temperatures were above normal parts of southern and eastern New Mexico.



Source: WestWide Drought Tracker

September – November precipitation totals were much-below normal for most of Arizona, and it was the driest September – November season on record for parts of western Arizona. In New Mexico, precipitation was much-above normal in some northern and eastern areas of the state, but in the southwest it ranged below to much-below normal.



Source: WestWide Drought Tracker

September – November temperatures were much-above normal across Arizona and New Mexico. In some areas it was the warmest September – November season on record.



Source: WestWide Drought Tracker

Drought

Drought conditions have not yet improved in Arizona and New Mexico. Drought or abnormally dry conditions are diagnosed for over half of New Mexico and about 90% of Arizona, by area. The most acute drought conditions are found in the southernmost parts of New Mexico and in western Arizona, where drought has been classified as D3 Extreme Drought. Areas of D1 Moderate Drought to D2 Severe Drought extend from those hardest-hit areas into central and southeast Arizona.



Source: U.S. Drought Monitor

NIDIS Improved and Expanded State Pages on Drought.Gov

Arizona

New Mexico

Hurricanes & Tropical Storms

The count of named storms in the eastern North Pacific for the 2024 hurricane season reached 12 in November with the addition of tropical storm Lane—fewer named storms than the long-term average of 15. Pacific tropical cyclones steered clear of the Southwest for the most part this year; arguably, the Atlantic had a greater impact with tropical storm Alberto bringing widespread rain to Arizona and New Mexico in late June.



Snowpack

Snowpack in Arizona and New Mexico is generally below normal to much-below normal. Snow water equivalent (SWE) is closer to normal in basins to the north—as of December 30, 2024, SWE aggregated for the Upper Colorado Basin is 95% of normal.



Water Supply

Water storage at Lake Mead and Lake Powell is well below long-term average levels, but close to where levels were last year. Other major reservoirs in Arizona are near or above long-term average levels, but in most cases holding less than last year. New Mexico reservoirs are generally below longterm average levels with a few exceptions—Ute Reservoir is near capacity, and Avalon is up over last year and above the long-term average.



The map gives a representation of current storage for reservoirs in Arizona and New Mexico. Reservoir locations are numbered within the blue circles on the map, corresponding to the reservoirs listed in the table. The cup next to each reservoir shows the current storage (blue fill) as a percent of total capacity. Note that while the size of each cup varies with the size of the reservoir, these are representational and not to scale. Each cup also represents last year's storage (dotted line) and the 1991–2020 reservoir average (red line). The table details more exactly the current capacity (listed as a percent of maximum storage). Current and maximum storage are given in thousands of acre-feet for each reservoir. One acre-foot is the volume of water sufficient to cover an acre of land to a depth of 1 foot (approximately 325,851 gallons). On average, 1 acre-foot of water is enough to meet the demands of four people for a year. The last column of the table lists an increase or decrease in storage since last month. A line indicates no change. These data are based on reservoir reports updated monthly by the <u>Natural</u> <u>Resources Conservation Service - National Water and Climate Center (USDA)</u>

BOR: New Mexico Dashboard

ENSO Tracker

November sea surface temperatures (SSTs) in the equatorial Pacific showed a La Niña-like pattern of below-normal SSTs in the central-eastern Pacific and above-normal SSTs in the west. But, as in the preceding months, the strength of the negative SST anomalies was not great enough to mark the beginning of a La Niña



Source: Australian Bureau of Meteorology

In the coming months, the La Niña-like pattern is forecast by most dynamical models to persist and strengthen, with the area of cooler-than normal SSTs crossing the threshold of 0.5°C below normal averaged over the Nino 3.4 monitoring region, as shown in this SST forecast from the Australian ACCESS forecast model.



Source: Australian Bureau of Meteorology

SSTs averaged for the ENSO monitoring regions have trended sharply toward cooler-than-normal temperatures over the month of December. By the latest weekly measurement, Nino 3.4 SST is 1.1°C below normal, which is cool enough to mark a transition to La Niña conditions.



Source: Climate Prediction Center (NOAA)

ENSO forecasts favor ENSO-neutral conditions for the coming months, assigning a probability of near 40% to La Niña conditions for the December – February forecast window, versus a near 60% probability of ENSO-neutral conditions. For subsequent forecast seasons ENSO-neutral is favored by increasingly larger probability margins, through Spring 2025.



The plume of individual ENSO model forecasts shows that ENSO-neutral conditions for the December – February forecast window are favored almost exclusively by statistical forecast models (those based on past ENSO behavior) while the dynamical models (those based on simulating physics) favor La Niña. Of those models that do predict La Niña, all show a peak in either the December – February or January – March forecast window and a relaxation toward ENSO-neutral in following months.

Source: The International Research Institute for Climate and Society, Columbia University Climate School

Seasonal Forecasts

The January – March seasonal precipitation forecast leans toward below normal precipitation (greater than 33% chance) for an area that includes Arizona and New Mexico. Southeastern Arizona and southern New Mexico are included in an area where the forecast calls below normal precipitation *likely*—with a probability between 50% and 60%.



Source: Climate Prediction Center (NOAA)

The January – March seasonal temperature forecast favors warmer-than-normal temperatures in Arizona and New Mexico. Both states are included in an area where the forecast either *leans* toward above normal temperatures (33% - 50% probability) or calls them *likely* (50% - 60% probability). The seasonal forecast generally follows the pattern of precipitation anomalies associated with La Niña; although by SST measures La Niña has been slow to emerge and apparently weak, the tropical atmosphere has been acting La Niña-like for several months, and it is the atmospheric response to ENSO that leads to seasonal climate anomalies in North America.



Source: Climate Prediction Center (NOAA)

Southwest Climate Podcast

December 2024 SW Climate Podcast - Relatively Speaking

Recorded 12/06/2024

Aired 12/10/2024

In this month's episode of the Southwest Climate Podcast hosts Zack Guido and Mike Crimmins slow-roll into the winter season. They look back on 2024 followed by what happened in November and why - including the Atmospheric River event on the west coast. They dive into a recent paper

that interrogates the expected intensification of cool season precipitation in the west. And close out with a teaser look at the La Niña outlook and precipitation forecast with a highlight of the Relative Oceanic Niño Index (RONI).

<u>Listen Here</u>

About CLIMAS

The Climate Assessment for the Southwest (CLIMAS) program was established in 1998 as part of the National Oceanic and Atmospheric Administration's Climate Adaptation Partnerships (CAP) Program (formerly known as Regional Integrated Sciences and Assessments, or RISA). CLIMAS housed at the University of Arizona's Institute of the Environment—is a collaboration between the University of Arizona and New Mexico State University. The CLIMAS team is made up of experts from a variety of social, physical, and natural sciences who work with partners across the Southwest to develop sustainable answers to regional climate challenges.



Learn more about the NOAA CAP program here

Disclaimer

This packet contains official and non-official forecasts, as well as other information. While we make every effort to verify this information, please understand that we do not warrant the accuracy of any of these materials. The user assumes the entire risk related to the use of this data. CLIMAS, and UA Cooperative Extension disclaim any and all warranties, whether expressed or implied, including (without limitation) any implied warranties of merchantability or fitness for a particular purpose. In no event will CLIMAS, UA Cooperative Extension, or The University of Arizona be liable to you or to any third party for any direct, indirect, incidental, consequential, special or exemplary damages or lost profit resulting from any use or misuse of this data.

> Southwest Climate Outlook contributors:

Mike Crimmins & Matt Meko